arXiv:math/9904047v5 [math.MG] 5 May 1999

Discrete Versions of the Beckman-Quarles Theorem

Apoloniusz Tyszka

to appear in Aequationes Mathematicae

Abstract. Let $\mathbf{F} \subseteq \mathbf{R}$ denote the field of numbers which are constructible by means of ruler and compass. We prove that: (1) if $x, y \in \mathbf{R}^n$ $(n > 1)$ and $|x - y|$ is an algebraic number then there exists a finite set $S_{xy} \subseteq \mathbf{R}^n$ containing x and y such that each map from S_{xy} to $\mathbf{R}^n$ preserving all unit distances preserves the distance between x and y; if $x, y \in \mathbf{F}^n$ then we can choose $S_{xy} \subseteq \mathbf{F}^n$, (2) only algebraic distances $|x - y|$ have the property from item (1), (3) if $X_1, X_2, ..., X_m \in \mathbf{R}^n$ $(n > 1)$ lie on some affine hyperplane then there exists a finite set $L(X_1, X_2, ..., X_m) \subseteq \mathbf{R}^n$ containing $X_1, X_2, ..., X_m$ such that each map from $L(X_1, X_2, ..., X_m)$ to $\mathbf{R}^n$ preserving all unit distances preserves the property that $X_1, X_2, ..., X_m$ lie on some affine hyperplane, (4) if $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |LM|$ $(|JK| < |LM|)$ then there exists a finite set $C_{JKLM} \subseteq \mathbf{R}^n$ containing J, K, L, M such that any map $f : C_{JKLM} \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(J)f(K)| = |f(L)f(M)|$ $(|f(J)f(K)| < |f(L)f(M)|)$.

Let $\mathbf{F} \subseteq \mathbf{R}$ denote the field of numbers which are constructible by means of ruler and compass. Theorem 1 may be viewed as a discrete form of the classical Beckman-Quarles theorem, which states that any map from $\mathbf{R}^n$ to $\mathbf{R}^n$ $(2 \leq n < \infty)$ preserving unit distances is an isometry, see [1], [2] and [5]. Theorem 1 was proved in [10] in the special case where $n = 2$ and the distance $|x - y|$ is constructible by means of ruler and compass.

Theorem 1. If $x, y \in \mathbf{R}^n$ $(n > 1)$ and $|x-y|$ is an algebraic number then there exists a finite set $S_{xy} \subseteq \mathbf{R}^n$ containing x and y such that each map from S_{xy} to $\mathbf{R}^n$ preserving all unit distances preserves the distance between x and y.

Proof. The proof falls naturally into two sections.

1. Elementary facts

Let us denote by D_n the set of all non-negative numbers d with the following property:

If $x, y \in \mathbf{R}^n$ and $|x-y| = d$ then there exists a finite set $S_{xy} \subseteq \mathbf{R}^n$ such that $x, y \in S_{xy}$ and any map $f : S_{xy} \to \mathbf{R}^n$ that preserves unit distance also preserves the distance between x and y.

Obviously $0, 1 \in D_n$. We first prove that if $d \in D_n$ then $\sqrt{2+2/n} \cdot d \in D_n$. Let us assume that $d > 0$, $x, y \in \mathbf{R}^n$, $|x-y| = \sqrt{2+2/n} \cdot d$. Using the notation of Figure 1 we show that

$$S_{xy} := \bigcup \{S_{ab} : a, b \in \{x, y, \widetilde{y}, p_1, p_2, ..., p_n, \widetilde{p}_1, \widetilde{p}_2, ..., \widetilde{p}_n\}, |a-b| = d\}$$

satisfies the condition of Theorem 1. Figure 1 shows the case $n = 2$, but equations below Figure 1 describe the general case $n \geq 2$; z denotes the centre of the $(n-1)$-dimensional regular simplex $p_1 p_2 ... p_n$.

Figure 1

$1 \leq i < j \leq n$

$|y - \widetilde{y}| = d$, $|x - p_i| = |y - p_i| = |p_i - p_j| = d = |x - \widetilde{p}_i| = |\widetilde{y} - \widetilde{p}_i| = |\widetilde{p}_i - \widetilde{p}_j|$

$|x - \widetilde{y}| = |x - y| = 2 \cdot |x - z| = 2 \cdot \sqrt{\frac{n+1}{2n}} \cdot d = \sqrt{2 + 2/n} \cdot d$

Let us assume that $f : S_{xy} \to \mathbf{R}^n$ preserves the distance 1. Since

$$S_{xy} \supseteq S_{y\widetilde{y}} \cup \bigcup_{i=1}^{n} S_{xp_i} \cup \bigcup_{i=1}^{n} S_{yp_i} \cup \bigcup_{1 \le i < j \le n} S_{p_i p_j}$$

we conclude that f preserves the distances between y and $\widetilde{y}$, x and p_i $(1 \le i \le n)$, y and p_i $(1 \le i \le n)$, and all distances between p_i and p_j $(1 \le i < j \le n)$. Hence $|f(y) - f(\widetilde{y})| = d$ and $|f(x) - f(y)|$ is either 0 or $\sqrt{2+2/n} \cdot d$. Analogously we have that $|f(x) - f(\widetilde{y})|$ is either 0 or $\sqrt{2+2/n} \cdot d$. Thus $f(x) \neq f(y)$, so $|f(x) - f(y)| = \sqrt{2+2/n} \cdot d$ which completes the proof that $\sqrt{2+2/n} \cdot d \in D_n$.

Hence, if $d \in D_n$ then $(2+2/n) \cdot d = \sqrt{2+2/n} \cdot (\sqrt{2+2/n} \cdot d) \in D_n$.

We next prove that if $x, y \in \mathbf{R}^n$, $d \in D_n$ and $|x - y| = (2/n) \cdot d$ then there exists a finite set $Z_{xy} \subseteq \mathbf{R}^n$ containing x and y such that any map $f : Z_{xy} \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(x) - f(y)| \le |x - y|$; this result is adapted from [5]. It is obvious if $n = 2$, therefore we assume that $n > 2$ and $d > 0$. Let us see at Figure 2 below, z denotes the centre of the $(n-1)$-dimensional regular simplex $p_1 p_2 ... p_n$. Figure 2 shows the case $n = 3$, but equations below Figure 2 describe the general case $n \ge 3$.

Figure 2

$$1 \le i < j \le n$$

$$|x - p_i| = |y - p_i| = d, \quad |p_i - p_j| = \sqrt{2+2/n} \cdot d, \quad |z - p_i| = \sqrt{1 - 1/n^2} \cdot d$$

$$|x - y| = 2 \cdot |x - z| = 2 \cdot \sqrt{|x - p_i|^2 - |z - p_i|^2} = 2 \cdot \sqrt{d^2 - (1 - 1/n^2) \cdot d^2} = (2/n) \cdot d$$

Let us define:

$$Z_{xy} := \bigcup_{1 \le i < j \le n} S_{p_i p_j} \cup \bigcup_{i=1}^{n} S_{x p_i} \cup \bigcup_{i=1}^{n} S_{y p_i}$$

If $f : Z_{xy} \to \mathbf{R}^n$ preserves the distance 1 then $|f(x) - f(y)| = |x - y| = (2/n) \cdot d$ or $|f(x) - f(y)| = 0$, hence $|f(x) - f(y)| \le |x - y|$.

If $d \in D_n$, then $2 \cdot d \in D_n$ (see Figure 3).

Figure 3

$|x - y| = 2 \cdot d$

$S_{xy} = S_{xs} \cup S_{sy} \cup Z_{yt} \cup S_{xt}$

From Figure 4 it is clear that if $d \in D_n$ then all distances $k \cdot d$ (k a positive integer) belong to D_n.

Figure 4

$|x - y| = k \cdot d$

$S_{xy} = \bigcup \{S_{ab} : a, b \in \{w_0, w_1, ..., w_k\}, |a - b| = d \vee |a - b| = 2 \cdot d\}$

From Figure 5 it is clear that if $d \in D_n$, then all distances d/k (k a positive integer) belong to D_n. Hence $D_n \supseteq Q^+$; here and subsequently Q^+ denotes the set of positive rational numbers.

Figure 5

$|x - y| = d/k$

$S_{xy} = S_{\tilde{x}\tilde{y}} \cup S_{\tilde{x}x} \cup S_{xz} \cup S_{\tilde{x}z} \cup S_{\tilde{y}y} \cup S_{yz} \cup S_{\tilde{y}z}$

Lemma. If $x, y \in \mathbf{R}^n$ $(n > 1)$ and $\varepsilon > 0$ then there exists a finite set $T_{xy}(\varepsilon) \subseteq \mathbf{R}^n$ containing x and y such that each map $f : T_{xy}(\varepsilon) \to \mathbf{R}^n$ preserving all unit distances preserves the distance between x and y to within ε i.e. $||f(x) - f(y)| - |x - y|| \le \varepsilon$.
Proof. It follows from Figure 6.

Figure 6
$|x - z|, |z - y| \in Q^+$, $|z - y| \le \varepsilon/2$
$T_{xy}(\varepsilon) = S_{xz} \cup S_{zy}$

2. Rigid graphs

Let $(\mathbf{R}^n, 1)$ denote the graph with vertex set $\mathbf{R}^n$ and with two vertices u, v adjacent if and only if $|u - v| = 1$. A *unit-distance graph* in $\mathbf{R}^n$ is any subgraph of $(\mathbf{R}^n, 1)$. A graph G in $\mathbf{R}^n$ with vertex set X is called *rigid* if there exists a $\delta > 0$ such that any $g : X \to \mathbf{R}^n$ satisfying: $|g(x) - x| < \delta$ for all $x \in X$ and $|g(x) - g(y)| = |x - y|$ for all edges xy of G, is an isometry on X.
H. Maehara proved that for any real algebraic number $r > 0$ there exists a finite rigid unit-distance graph G in $\mathbf{R}^n$ $(n > 1)$ having two vertices α and β at distance r, see [7] for $n = 2$ and [7]-[8] for $n > 2$. Let us denote by X the set of vertices of G. Let us fix: $\delta > 0$ from the definition of rigidity and $p_1, ..., p_n, p_{n+1} \in \mathbf{R}^n$ such that $|p_i - p_j| = 1$ for $1 \le i < j \le n + 1$. Since $p_1, ..., p_n, p_{n+1}$ are affinely independent, the following mapping

$$\mathbf{R}^n \ni y \xrightarrow{\Phi} (|p_1 - y|, ..., |p_n - y|, |p_{n+1} - y|) \in \mathbf{R}^{n+1}$$

is injective, see [3] p.127.

Φ is continuous, X is finite. We can choose $\varepsilon > 0$ such that for all $x \in X$, $\widetilde{x} \in \mathbf{R}^n$

$$max_{1 \le i \le n+1} ||p_i - \widetilde{x}| - |p_i - x|| \le \varepsilon \text{ implies } |\widetilde{x} - x| < \delta.$$

We prove that $S_{\alpha\beta} := \bigcup_{x \in X} \bigcup_{i=1}^{n+1} T_{p_i x}(\varepsilon)$ satisfies the condition of the theorem. Let us assume that $f : S_{\alpha\beta} \to \mathbf{R}^n$ preserves the distance 1. Hence f preserves distances between p_i and p_j for $1 \le i < j \le n+1$. There is an isometry $I : \mathbf{R}^n \to \mathbf{R}^n$ such that $f(p_i) = I(p_i)$ for $1 \le i \le n+1$. Since $S_{\alpha\beta} \supseteq T_{p_i x}(\varepsilon)$ $(1 \le i \le n+1)$, f preserves distances between p_i and $x \in X$ to within ε. Also $I^{-1} \circ f$ preserves distances between p_i and $x \in X$ to within ε. Therefore, for all $x \in X$

$$max_{1 \le i \le n+1} ||(I^{-1} \circ f)(p_i) - (I^{-1} \circ f)(x)| - |p_i - x|| \le \varepsilon$$

i.e.

$$max_{1 \le i \le n+1} ||p_i - (I^{-1} \circ f)(x)| - |p_i - x|| \le \varepsilon$$

Hence, for all $x \in X$ $|(I^{-1} \circ f)(x) - x| < \delta$. If xy is an edge of G, then $|x - y| = 1$. Hence $|(I^{-1} \circ f)(x) - (I^{-1} \circ f)(y)| = 1 = |x - y|$. By the rigidity of G, $I^{-1} \circ f$ is an isometry on X. In particular, $I^{-1} \circ f$ preserves the distance between α and β. Hence f preserves the distance between α and β, which ends the proof.

Remark 1. The lemma from the proof of Theorem 1 implies the classical Beckman-Quarles theorem.

Let $L = (+, \cdot, \le, 0, 1)$ denote the language of ordered fields. We say that $X \subseteq \mathbf{R}^n$ is definable using L if there is an L-formula $\phi(x_1, ..., x_n)$ such that $X = \{(a_1, ..., a_n) \in \mathbf{R}^n : \phi(a_1, ..., a_n) \ holds\}$

Remark 2. Only algebraic distances $|x - y|$ have the property from Theorem 1. Indeed, if $x_1, x_2 \in \mathbf{R}^n$ $(n > 1)$ and $S_{x_1 x_2} = \{x_1, x_2, ..., x_m\}$ then $\{|x_1 - x_2|\} \subseteq \mathbf{R}$ is definable using L by the following L-formula $\psi(z)$:

$$(0 \le z) \wedge \forall x_{11} ... \forall x_{1n} ... \forall x_{m1} ... \forall x_{mn} (\underbrace{... \wedge (x_{i1} - x_{j1})^2 + ... + (x_{in} - x_{jn})^2 = 1 \wedge ...}_{(i,j) \in \{(i,j) : 1 \le i < j \le m, |x_i - x_j| = 1\}}$$

$$\Rightarrow (x_{11} - x_{21})^2 + ... + (x_{1n} - x_{2n})^2 = z^2)$$

From this, we deduce that $|x_1 - x_2|$ is an algebraic number defined by some quantifier free L-formula equivalent to $\psi(z)$, see [4].

Remark 3. M. Koshelev proved that Theorem 1 remains valid for rational distances in two-dimensional Minkowski space-time, see [6].

Theorem 2. If $x, y \in \mathbf{F}^n$ $(n > 1)$ then there exists a finite set $S_{xy} \subseteq \mathbf{F}^n$ containing x and y such that each map from S_{xy} to $\mathbf{R}^n$ preserving all unit distances preserves the distance between x and y.

Proof. Let us denote by C_n $(n > 1)$ the set of all non-negative constructible numbers d with the following property:

If $x, y \in \mathbf{F}^n$ and $|x - y| = d$ then there exists a finite set $S_{xy} \subseteq \mathbf{F}^n$ such that $x, y \in S_{xy}$ and any map $f : S_{xy} \to \mathbf{R}^n$ that preserves unit distance also preserves the distance between x and y.

Obviously $0, 1 \in C_n$. Similarly as in the proof of Theorem 1 we can prove that:

1) $\sqrt{2 + 2/n} \in C_n$,

2) if $d \in C_n$ then all distances $r \cdot d$ (r is positive and rational) belong to C_n, therefore all non-negative rational numbers belong to C_n,

3) if $x, y \in \mathbf{F}^n$ $(n > 1)$ and $\varepsilon > 0$ then there exists a finite set $T_{xy}(\varepsilon) \subseteq \mathbf{F}^n$ containing x and y such that each map $f : T_{xy}(\varepsilon) \to \mathbf{R}^n$ preserving all unit distances preserves the distance between x and y to within ε i.e. $||f(x) - f(y)| - |x - y|| \leq \varepsilon$.

If $a, b \in C_n$, $a > b > 0$ then $\sqrt{a^2 - b^2} \in C_n$ (see Figure 7).

Figure 7
$|x - y| = \sqrt{a^2 - b^2}$
$S_{xy} = S_{sx} \cup S_{xt} \cup S_{st} \cup S_{sy} \cup S_{ty}$

Hence $\sqrt{3} \cdot a = \sqrt{(2 \cdot a)^2 - a^2} \in C_n$ and $\sqrt{2} \cdot a = \sqrt{(\sqrt{3} \cdot a)^2 - a^2} \in C_n$. Therefore $\sqrt{a^2 + b^2} = \sqrt{(\sqrt{2} \cdot a)^2 - (\sqrt{a^2 - b^2})^2} \in C_n$.

Let us see at Figure 8 below, z denotes the centre of the $(n-1)$-dimensional regular simplex $p_1 p_2 ... p_n$, $n = 2$. This construction shows that if $a, b \in C_n$, $a > b > 0$, $n \geq 2$ (consistently, equations below Figure 8 describe the general case $n \geq 2$) then $a - b \in C_n$, hence $a + b = 2 \cdot a - (a - b) \in C_n$.

Figure 8

$$|x-y| = a-b, \;\; |x-z| = a \in C_n, \;\; |y-z| = b \in C_n$$

$$|p_i - p_j| = \sqrt{2 + 2/n} \in C_n, \;\; |z - p_i| = \sqrt{1^2 - (1/n)^2} \in C_n, \;\; 1 \leq i < j \leq n$$

$$|x - p_1| = \sqrt{|x-z|^2 + |z-p_1|^2} = ... = |x - p_n| = \sqrt{|x-z|^2 + |z-p_n|^2} \in C_n$$

$$|y - p_1| = \sqrt{|y-z|^2 + |z-p_1|^2} = ... = |y - p_n| = \sqrt{|y-z|^2 + |z-p_n|^2} \in C_n$$

$$S_{xy} = \bigcup_{1 \leq i < j \leq n} S_{p_i p_j} \cup \bigcup_{i=1}^{n} S_{x p_i} \cup \bigcup_{i=1}^{n} S_{y p_i} \cup T_{xy}(b)$$

In order to prove that $C_n \setminus \{0\}$ is a multiplicative group it remains to observe that if positive $a, b, c \in C_n$ then $\frac{a \cdot b}{c} \in C_n$ (see Figure 9).

Figure 9

m is positive and integer

$$b < 2 \cdot m \cdot c$$

$$S_{AB} = S_{OA} \cup S_{OB} \cup S_{O\widetilde{A}} \cup S_{O\widetilde{B}} \cup S_{A\widetilde{A}} \cup S_{B\widetilde{B}} \cup S_{\widetilde{A}\widetilde{B}}$$

If $a \in C_n$, $a > 1$, then $\sqrt{a} = \frac{1}{2} \cdot \sqrt{(a+1)^2 - (a-1)^2} \in C_n$; if $a \in C_n$, $0 < a < 1$, then $\sqrt{a} = 1/\sqrt{\frac{1}{a}} \in C_n$. Thus C_n contains all non-negative real numbers contained in the real quadratic closure of Q. This completes the proof.

Directly from the Lemma (see the proof of Theorem 1) we obtain the following two corollaries, these corollaries and the next elementary proposition are useful in our next proofs.

Corollary 1. If $p \neq q \in \mathbf{R}^n$ $(n > 1)$ then any map $f : T_{pq}(|pq|/2) \to \mathbf{R}^n$ that preserves unit distance satisfies $f(p) \neq f(q)$.

Corollary 2. If $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| < |LM|$ then any map $f : T_{JK}(\frac{|LM|-|JK|}{3}) \cup T_{LM}(\frac{|LM|-|JK|}{3}) \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(J)f(K)| < |f(L)f(M)|$.

Proposition. If $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |LM|$ then there exist $A, B \in \mathbf{R}^n$ such that:
(1) segments JK and AB are symmetric with respect to some hyperplane $H(JK, AB)$
and
(2) segments AB and LM are symmetric with respect to some hyperplane $H(AB, LM)$.
Proof. Throughout the proof for any two points $x \in \mathbf{R}^n$ and $y \in \mathbf{R}^n$ H_{xy} stands for the hyperplane which reflects x into y, H_{xy} is unique if $x \neq y$. Let $\widetilde{H}_{xy} : \mathbf{R}^n \to \mathbf{R}^n$ denote the hyperplane symmetry determined by H_{xy}. We put: $A := L = \widetilde{H}_{JL}(J)$, $B := \widetilde{H}_{JL}(K)$, $H(JK, AB) := H_{JL}$. Since $|AB| = |JK| = |LM| = |AM|$ there exists a hyperplane H_{BM} satisfying $\widetilde{H}_{BM}(A) = A = L$. Therefore, we put $H(AB, LM) := H_{BM}$, obviously $\widetilde{H}(AB, LM)(B) = M$. This completes the proof.

Theorems 3 and 4 below yield information about unit-distance preserving mappings on finite subsets of $\mathbf{R}^n$. We present their proofs omitting details.

Theorem 3. If $X_1, X_2, ..., X_m \in \mathbf{R}^n$ $(n > 1)$ lie on some affine hyperplane then there exists a finite set $L(X_1, X_2, ..., X_m) \subseteq \mathbf{R}^n$ containing $X_1, X_2, ..., X_m$ such that each map from $L(X_1, X_2, ..., X_m)$ to $\mathbf{R}^n$ preserving all unit distances preserves the property that $X_1, X_2, ..., X_m$ lie on some affine hyperplane. If $X_1, X_2, ..., X_m \in \mathbf{F}^n$ then we can choose $L(X_1, X_2, ..., X_m) \subseteq \mathbf{F}^n$.

Proof. We choose an affine hyperplane G such that $X_1, X_2, ..., X_m \in G$. We fix $P \in G$ and choose an integer u such that $|PX_i| \leq 2 \cdot u$ for $1 \leq i \leq m$. We fix $O \in \mathbf{R}^n$ such that PO is perpendicular to G and $|PO| = 4 \cdot u$. Let us see at Figure 10 below, this drawing shows the case $n = 2$, but equations below Figure 10 describe the general case $n \geq 2$.

Figure 10

$$\forall i \in \{1, 2, ..., m\} \forall j \in \{1, 2, ..., n\} |X_i B_{ij}| = |A_i B_{ij}| = 2 \cdot u \wedge |OB_{ij}| = 6 \cdot u$$

$$|PO| = 4 \cdot u, \ \forall i \in \{1, 2, ..., m\} |PA_i| = 4 \cdot u \wedge |PT_i| = |T_i X_i| = u$$

Erasing all dashed lines and segments PT_i, $T_i X_i$, PA_i, PO from Figure 10 we obtain the Peaucellier inversor:$|OX_i| \cdot |OA_i| = |OC_i| \cdot |OD_i| = (6 \cdot u - 2 \cdot u) \cdot (6 \cdot u + 2 \cdot u) = (6 \cdot u)^2 - (2 \cdot u)^2$ (see [9] in the case $n = 2$). Always:

$|PX_i| \le 2 \cdot u < |PA_i|$. Therefore, if all edge-lengths (continuous segments) are preserved and if all relations $B_{ik} \ne B_{il}$ $(1 \le k < l \le n)$ are preserved then PX_i is perpendicular to PO. From this, we conclude that

$$L_i := S_{PO} \cup S_{PT_i} \cup S_{T_iX_i} \cup S_{PA_i} \cup (\bigcup_{j=1}^{n} S_{X_iB_{ij}} \cup S_{A_iB_{ij}} \cup S_{OB_{ij}}) \cup$$
$$\bigcup \{T_{B_{ik}B_{il}}(|B_{ik}B_{il}|/2) : 1 \le k < l \le n\}$$

is adequate for preserving the property that PX_i is perpendicular to PO. Hence $L(X_1, X_2, ..., X_m) := \bigcup_{i=1}^{m} L_i$ satisfies the condition of the theorem. This completes the proof for $\mathbf{R}^n$. The proof for $\mathbf{F}^n$ is similar.

Theorem 4 (cf. Corollary 2). If $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |LM|$ then there exists a finite set $C_{JKLM} \subseteq \mathbf{R}^n$ containing J, K, L, M such that any map $f : C_{JKLM} \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(J)f(K)| = |f(L)f(M)|$. If $J, K, L, M \in \mathbf{F}^n$ then we can choose $C_{JKLM} \subseteq \mathbf{F}^n$.

Proof. Let us observe that if $J, K, A, B, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |AB| = |LM|$ then $C_{JKLM} := C_{JKAB} \cup C_{ABLM}$ satisfies the condition of Theorem 4. By Proposition it is sufficient to prove Theorem 4 in the case where two segments of equal lengths are symmetric with respect to some hyperplane H.

Let us see at figures below, drawings show the case $n = 2$, but equations below figures describe the general case $n \ge 2$.

Figure 11a

s, t are positive and integer

$$\forall j \in \{1, 2, ..., n\} |X_jJ| = |X_jL| = s \wedge |Y_jK| = |Y_jM| = t$$

$$|JK| = |LM| < |JM| = |LK| \qquad \delta := \frac{|JM| - |JK|}{3}$$

Figure 11b

s, t are positive and integer

$\forall j \in \{1, 2, ..., n\} |X_j J| = |X_j L| = s \wedge |Y_j K| = |Y_j M| = t$

$|JK| = |LM| > |JM| = |LK| \qquad \delta := \frac{|JK| - |JM|}{3}$

If all edge-lengths (continuous segments) are preserved and if all relations: $J \neq L$,$X_k \neq X_l$,$K \neq M$,$Y_k \neq Y_l$ $(1 \leq k < l \leq n)$ are preserved then images of J and L are symmetric and images of K and M are symmetric. Hence: $C_{JKLM} := L(X_1, X_2, ..., X_n, Y_1, Y_2, ..., Y_n) \cup (\bigcup_{j=1}^{n} S_{X_j J} \cup S_{X_j L}) \cup T_{JL}(|JL|/2) \cup \bigcup \{T_{X_k X_l}(|X_k X_l|/2) : 1 \leq k < l \leq n\} \cup (\bigcup_{j=1}^{n} S_{Y_j K} \cup S_{Y_j M}) \cup T_{KM}(|KM|/2) \cup \bigcup \{T_{Y_k Y_l}(|Y_k Y_l|/2) : 1 \leq k < l \leq n\} \cup T_{JK}(\delta) \cup T_{LM}(\delta) \cup T_{JM}(\delta) \cup T_{LK}(\delta)$ satisfies the condition of the theorem.

Separate formulas for C_{JKLM} are necessary if $J = L$ or $K = M$. If $J = L$ and $K = M$ then $C_{JKLM} := \{J, K\}$, if $J \neq L$ and $K = M$ then we see at Figure 11c, Figure 11c shows the case $n = 2$, but equations below Figure 11c describe the general case $n \geq 2$.

Figure 11c

s is positive and integer, $\qquad \forall j \in \{1, 2, ..., n\} |X_j J| = |X_j L| = s$

$C_{JKLM} = L(X_1, X_2, ..., X_n, K) \cup (\bigcup_{j=1}^{n} S_{X_j J} \cup S_{X_j L}) \cup T_{JL}(|JL|/2) \cup$

$\bigcup \{T_{X_k X_l}(|X_k X_l|/2) : 1 \leq k < l \leq n\}$

Symmetric construction applies to the case: $J = L$ and $K \neq M$. All cases are explained and the proof for $\mathbf{R}^n$ is complete. The proof for $\mathbf{F}^n$ is similar.

Conjecture. Let us assume that $X \subseteq \underbrace{\mathbf{R}^n \times ... \times \mathbf{R}^n}_{m-times} \cong \mathbf{R}^{mn}$ $(n > 1, m \geq 1)$ is definable using L and the relation X is preserved under isometries of $\mathbf{R}^n$. We suppose that if $(x_1, ..., x_m) \in X$ then there exists a finite set $P(x_1, ..., x_m) \subseteq \mathbf{R}^n$ containing $x_1, ..., x_m$ such that any map $f : P(x_1, ..., x_m) \to \mathbf{R}^n$ that preserves unit distance satisfies $(f(x_1), ..., f(x_m)) \in X$.

References

1. F. S. Beckman and D. A. Quarles Jr., On isometries of euclidean spaces, *Proc. Amer. Math. Soc.*, 4 (1953), 810-815.

2. W. Benz, *Geometrische Transformationen (unter besonderer Berücksichtigung der Lorentztransformationen).* BI Wissenschaftsverlag, Mannheim, Leipzig, Wien, Zürich, 1992.

3. K. Borsuk, *Multidimensional analytic geometry*, Polish Scientific Publishers, Warsaw, 1969.

4. C. C. Chang and H. J. Keisler, *Model Theory*, 3rd ed., North-Holland, Amsterdam, 1990.

5. U. Everling, Solution of the isometry problem stated by K. Ciesielski, *Math. Intelligencer*, 10 (1988), No.4, p.47.

6. M. Koshelev, We can measure any distance or any amount of time with a most primitive clock and a most primitive ruler: a space-time version of Tyszka's result, *Geombinatorics*, 7 (1998), No.3, 95-100.

7. H. Maehara, Distances in a rigid unit-distance graph in the plane, *Discrete Appl. Math.* 31 (1991), 193-200.

8. H. Maehara, Extending a flexible unit-bar framework to a rigid one, *Discrete Math.* 108 (1992), 167-174.

9. H. Rademacher and O. Toeplitz, *The enjoyment of mathematics*, Princeton University Press, Princeton, 1994.
10. A. Tyszka, A discrete form of the Beckman-Quarles theorem, *Amer. Math. Monthly* 104 (1997), 757-761.

Technical Faculty
Hugo Kołłątaj University
Balicka 104, PL-30-149 Kraków, Poland
rttyszka@cyf-kr.edu.pl
http://www.cyf-kr.edu.pl/~rttyszka

This figure "figure1.gif" is available in "gif" format from:

This figure "figure2.gif" is available in "gif" format from:

This figure "figure3.gif" is available in "gif" format from:

This figure "figure4.gif" is available in "gif" format from:

This figure "figure5.gif" is available in "gif" format from:

This figure "figure6.gif" is available in "gif" format from:

This figure "figure7.gif" is available in "gif" format from:

This figure "figure8.gif" is available in "gif" format from:

This figure "figure9.gif" is available in "gif" format from:

This figure "figure10.gif" is available in "gif" format from:

http://arXiv.org/ps/math/9904047v5

This figure "figur11a.gif" is available in "gif" format from:

This figure "figur11b.gif" is available in "gif" format from:

This figure "figur11c.gif" is available in "gif" format from:

Discrete versions of the Beckman-Quarles theorem

Apoloniusz Tyszka

to appear in Aequationes Mathematicae

Abstract. Let $\mathbf{F} \subseteq \mathbf{R}$ denote the field of numbers which are constructible by means of ruler and compass. We prove that: (1) if $x, y \in \mathbf{R}^n$ $(n > 1)$ and $|x - y|$ is an algebraic number then there exists a finite set $S_{xy} \subseteq \mathbf{R}^n$ containing x and y such that each map from S_{xy} to $\mathbf{R}^n$ preserving all unit distances preserves the distance between x and y; if $x, y \in \mathbf{F}^n$ then we can choose $S_{xy} \subseteq \mathbf{F}^n$, (2) only algebraic distances $|x - y|$ have the property from item (1), (3) if $X_1, X_2, ..., X_m \in \mathbf{R}^n$ $(n > 1)$ lie on some affine hyperplane then there exists a finite set $L(X_1, X_2, ..., X_m) \subseteq \mathbf{R}^n$ containing $X_1, X_2, ..., X_m$ such that each map from $L(X_1, X_2, ..., X_m)$ to $\mathbf{R}^n$ preserving all unit distances preserves the property that $X_1, X_2, ..., X_m$ lie on some affine hyperplane, (4) if $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |LM|$ $(|JK| < |LM|)$ then there exists a finite set $C_{JKLM} \subseteq \mathbf{R}^n$ containing J, K, L, M such that any map $f : C_{JKLM} \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(J)f(K)| = |f(L)f(M)|$ $(|f(J)f(K)| < |f(L)f(M)|)$.

Let $\mathbf{F} \subseteq \mathbf{R}$ denote the field of numbers which are constructible by means of ruler and compass. Theorem 1 may be viewed as a discrete form of the classical Beckman-Quarles theorem, which states that any map from $\mathbf{R}^n$ to $\mathbf{R}^n$ $(2 \leq n < \infty)$ preserving unit distances is an isometry, see [1], [2] and [5]. Theorem 1 was proved in [10] in the special case where $n = 2$ and the distance $|x - y|$ is constructible by means of ruler and compass.

Theorem 1. If $x, y \in \mathbf{R}^n$ $(n > 1)$ and $|x-y|$ is an algebraic number then there exists a finite set $S_{xy} \subseteq \mathbf{R}^n$ containing x and y such that each map from S_{xy} to $\mathbf{R}^n$ preserving all unit distances preserves the distance between x and y.

Proof. The proof falls naturally into two sections.

1. Elementary facts

Let us denote by D_n the set of all non-negative numbers d with the following property:

If $x, y \in \mathbf{R}^n$ and $|x-y| = d$ then there exists a finite set $S_{xy} \subseteq \mathbf{R}^n$ such that $x, y \in S_{xy}$ and any map $f : S_{xy} \to \mathbf{R}^n$ that preserves unit distance also preserves the distance between x and y.

Obviously $0, 1 \in D_n$. We first prove that if $d \in D_n$ then $\sqrt{2+2/n} \cdot d \in D_n$. Let us assume that $d > 0$, $x, y \in \mathbf{R}^n$, $|x-y| = \sqrt{2+2/n} \cdot d$. Using the notation of Figure 1 we show that

$$S_{xy} := \bigcup \{S_{ab} : a, b \in \{x, y, \tilde{y}, p_1, p_2, ..., p_n, \tilde{p}_1, \tilde{p}_2, ..., \tilde{p}_n\}, |a-b| = d\}$$

satisfies the condition of Theorem 1. Figure 1 shows the case $n = 2$, but equations below Figure 1 describe the general case $n \geq 2$; z denotes the centre of the $(n-1)$-dimensional regular simplex $p_1 p_2 ... p_n$.



Figure 1

$$1 \leq i < j \leq n$$

$$|y-\tilde{y}| = d,\ |x-p_i| = |y-p_i| = |p_i-p_j| = d = |x-\tilde{p}_i| = |\tilde{y}-\tilde{p}_i| = |\tilde{p}_i-\tilde{p}_j|$$

$$|x-\tilde{y}| = |x-y| = 2 \cdot |x-z| = 2 \cdot \sqrt{\tfrac{n+1}{2n}} \cdot d = \sqrt{2+2/n} \cdot d$$

Let us assume that $f : S_{xy} \to \mathbf{R}^n$ preserves the distance 1. Since

$$S_{xy} \supseteq S_{y\widetilde{y}} \cup \bigcup_{i=1}^{n} S_{xp_i} \cup \bigcup_{i=1}^{n} S_{yp_i} \cup \bigcup_{1\le i<j\le n} S_{p_ip_j}$$

we conclude that f preserves the distances between y and $\widetilde{y}$, x and p_i $(1 \le i \le n)$, y and p_i $(1 \le i \le n)$, and all distances between p_i and p_j $(1 \le i < j \le n)$. Hence $|f(y) - f(\widetilde{y})| = d$ and $|f(x) - f(y)|$ is either 0 or $\sqrt{2+2/n}\cdot d$. Analogously we have that $|f(x) - f(\widetilde{y})|$ is either 0 or $\sqrt{2+2/n}\cdot d$. Thus $f(x) \ne f(y)$, so $|f(x) - f(y)| = \sqrt{2+2/n}\cdot d$ which completes the proof that $\sqrt{2+2/n}\cdot d \in D_n$.

Hence, if $d \in D_n$ then $(2+2/n)\cdot d = \sqrt{2+2/n}\cdot(\sqrt{2+2/n}\cdot d) \in D_n$.

We next prove that if $x, y \in \mathbf{R}^n$, $d \in D_n$ and $|x-y| = (2/n)\cdot d$ then there exists a finite set $Z_{xy} \subseteq \mathbf{R}^n$ containing x and y such that any map $f : Z_{xy} \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(x) - f(y)| \le |x-y|$; this result is adapted from [5]. It is obvious if $n = 2$, therefore we assume that $n > 2$ and $d > 0$. Let us see at Figure 2 below, z denotes the centre of the $(n-1)$-dimensional regular simplex $p_1p_2...p_n$. Figure 2 shows the case $n = 3$, but equations below Figure 2 describe the general case $n \ge 3$.



Figure 2

$$1 \le i < j \le n$$

$$|x-p_i| = |y-p_i| = d, \quad |p_i - p_j| = \sqrt{2+2/n}\cdot d, \quad |z-p_i| = \sqrt{1-1/n^2}\cdot d$$

$$|x-y| = 2\cdot|x-z| = 2\cdot\sqrt{|x-p_i|^2 - |z-p_i|^2} = 2\cdot\sqrt{d^2 - (1-1/n^2)\cdot d^2} = (2/n)\cdot d$$

Let us define:

$$Z_{xy} := \bigcup_{1 \le i < j \le n} S_{p_i p_j} \cup \bigcup_{i=1}^{n} S_{x p_i} \cup \bigcup_{i=1}^{n} S_{y p_i}$$

If $f : Z_{xy} \to \mathbf{R}^n$ preserves the distance 1 then $|f(x) - f(y)| = |x - y| = (2/n) \cdot d$ or $|f(x) - f(y)| = 0$, hence $|f(x) - f(y)| \le |x - y|$.

If $d \in D_n$, then $2 \cdot d \in D_n$ (see Figure 3).



Figure 3

$|x - y| = 2 \cdot d$

$S_{xy} = S_{xs} \cup S_{sy} \cup Z_{yt} \cup S_{xt}$

From Figure 4 it is clear that if $d \in D_n$ then all distances $k \cdot d$ (k a positive integer) belong to D_n.



Figure 4

$|x - y| = k \cdot d$

$S_{xy} = \bigcup\{S_{ab} : a, b \in \{w_0, w_1, ..., w_k\}, |a - b| = d \vee |a - b| = 2 \cdot d\}$

From Figure 5 it is clear that if $d \in D_n$, then all distances d/k (k a positive integer) belong to D_n. Hence $D_n \supseteq Q^+$; here and subsequently Q^+ denotes the set of positive rational numbers.



Figure 5

$|x - y| = d/k$

$S_{xy} = S_{\tilde{x}\tilde{y}} \cup S_{\tilde{x}x} \cup S_{xz} \cup S_{\tilde{x}z} \cup S_{\tilde{y}y} \cup S_{yz} \cup S_{\tilde{y}z}$

Lemma. If $x, y \in \mathbf{R}^n$ $(n > 1)$ and $\varepsilon > 0$ then there exists a finite set $T_{xy}(\varepsilon) \subseteq \mathbf{R}^n$ containing x and y such that each map $f : T_{xy}(\varepsilon) \to \mathbf{R}^n$ preserving all unit distances preserves the distance between x and y to within ε i.e. $||f(x) - f(y)| - |x - y|| \le \varepsilon$.
Proof. It follows from Figure 6.



Figure 6
$|x - z|, |z - y| \in Q^+$, $|z - y| \le \varepsilon/2$
$T_{xy}(\varepsilon) = S_{xz} \cup S_{zy}$

2. Rigid graphs

Let $(\mathbf{R}^n, 1)$ denote the graph with vertex set $\mathbf{R}^n$ and with two vertices u, v adjacent if and only if $|u - v| = 1$. A *unit-distance graph* in $\mathbf{R}^n$ is any subgraph of $(\mathbf{R}^n, 1)$. A graph G in $\mathbf{R}^n$ with vertex set X is called *rigid* if there exists a $\delta > 0$ such that any $g : X \to \mathbf{R}^n$ satisfying: $|g(x) - x| < \delta$ for all $x \in X$ and $|g(x) - g(y)| = |x - y|$ for all edges xy of G, is an isometry on X.
H. Maehara proved that for any real algebraic number $r > 0$ there exists a finite rigid unit-distance graph G in $\mathbf{R}^n$ $(n > 1)$ having two vertices α and β at distance r, see [7] for $n = 2$ and [7]-[8] for $n > 2$. Let us denote by X the set of vertices of G. Let us fix: $\delta > 0$ from the definition of rigidity and $p_1, ..., p_n, p_{n+1} \in \mathbf{R}^n$ such that $|p_i - p_j| = 1$ for $1 \le i < j \le n + 1$. Since $p_1, ..., p_n, p_{n+1}$ are affinely independent, the following mapping

$$\mathbf{R}^n \ni y \xrightarrow{\Phi} (|p_1 - y|, ..., |p_n - y|, |p_{n+1} - y|) \in \mathbf{R}^{n+1}$$

is injective, see [3] p.127.

Φ is continuous, X is finite. We can choose $\varepsilon > 0$ such that for all $x \in X$, $\widetilde{x} \in \mathbf{R}^n$

$$max_{1 \le i \le n+1} ||p_i - \widetilde{x}| - |p_i - x|| \le \varepsilon \text{ implies } |\widetilde{x} - x| < \delta.$$

We prove that $S_{\alpha\beta} := \bigcup_{x\in X} \bigcup_{i=1}^{n+1} T_{p_i x}(\varepsilon)$ satisfies the condition of the theorem. Let us assume that $f : S_{\alpha\beta} \to \mathbf{R}^n$ preserves the distance 1. Hence f preserves distances between p_i and p_j for $1 \le i < j \le n+1$. There is an isometry $I : \mathbf{R}^n \to \mathbf{R}^n$ such that $f(p_i) = I(p_i)$ for $1 \le i \le n+1$. Since $S_{\alpha\beta} \supseteq T_{p_i x}(\varepsilon)$ $(1 \le i \le n+1)$, f preserves distances between p_i and $x \in X$ to within ε. Also $I^{-1} \circ f$ preserves distances between p_i and $x \in X$ to within ε. Therefore, for all $x \in X$

$$max_{1\le i\le n+1} ||(I^{-1}\circ f)(p_i) - (I^{-1}\circ f)(x)| - |p_i - x|| \le \varepsilon$$

i.e.

$$max_{1\le i\le n+1} ||p_i - (I^{-1}\circ f)(x)| - |p_i - x|| \le \varepsilon$$

Hence, for all $x \in X$ $|(I^{-1}\circ f)(x) - x| < \delta$. If xy is an edge of G, then $|x-y| = 1$. Hence $|(I^{-1}\circ f)(x) - (I^{-1}\circ f)(y)| = 1 = |x-y|$. By the rigidity of G, $I^{-1}\circ f$ is an isometry on X. In particular, $I^{-1}\circ f$ preserves the distance between α and β. Hence f preserves the distance between α and β, which ends the proof.

Remark 1. The lemma from the proof of Theorem 1 implies the classical Beckman-Quarles theorem.

Let $L = (+, \cdot, \le, 0, 1)$ denote the language of ordered fields. We say that $X \subseteq \mathbf{R}^n$ is definable using L if there is an L-formula $\phi(x_1, ..., x_n)$ such that $X = \{(a_1, ..., a_n) \in \mathbf{R}^n : \phi(a_1, ..., a_n)\ holds\}$

Remark 2. Only algebraic distances $|x-y|$ have the property from Theorem 1. Indeed, if $x_1, x_2 \in \mathbf{R}^n$ $(n > 1)$ and $S_{x_1 x_2} = \{x_1, x_2, ..., x_m\}$ then $\{|x_1 - x_2|\} \subseteq \mathbf{R}$ is definable using L by the following L-formula $\psi(z)$:

$$(0 \le z) \wedge \forall x_{11} ... \forall x_{1n} ... \forall x_{m1} ... \forall x_{mn} (\underbrace{... \wedge (x_{i1} - x_{j1})^2 + ... + (x_{in} - x_{jn})^2 = 1 \wedge ...}_{(i,j)\in\{(i,j):1\le i<j\le m, |x_i - x_j| = 1\}}$$

$$\Rightarrow (x_{11} - x_{21})^2 + ... + (x_{1n} - x_{2n})^2 = z^2)$$

From this, we deduce that $|x_1 - x_2|$ is an algebraic number defined by some quantifier free L-formula equivalent to $\psi(z)$, see [4].

Remark 3. M. Koshelev proved that Theorem 1 remains valid for rational distances in two-dimensional Minkowski space-time, see [6].

Theorem 2. If $x, y \in \mathbf{F}^n$ $(n > 1)$ then there exists a finite set $S_{xy} \subseteq \mathbf{F}^n$ containing x and y such that each map from S_{xy} to $\mathbf{R}^n$ preserving all unit distances preserves the distance between x and y.

Proof. Let us denote by C_n $(n > 1)$ the set of all non-negative constructible numbers d with the following property:

If $x, y \in \mathbf{F}^n$ and $|x - y| = d$ then there exists a finite set $S_{xy} \subseteq \mathbf{F}^n$ such that $x, y \in S_{xy}$ and any map $f : S_{xy} \to \mathbf{R}^n$ that preserves unit distance also preserves the distance between x and y.

Obviously $0, 1 \in C_n$. Similarly as in the proof of Theorem 1 we can prove that:

1) $\sqrt{2 + 2/n} \in C_n$,

2) if $d \in C_n$ then all distances $r \cdot d$ (r is positive and rational) belong to C_n, therefore all non-negative rational numbers belong to C_n,

3) if $x, y \in \mathbf{F}^n$ $(n > 1)$ and $\varepsilon > 0$ then there exists a finite set $T_{xy}(\varepsilon) \subseteq \mathbf{F}^n$ containing x and y such that each map $f : T_{xy}(\varepsilon) \to \mathbf{R}^n$ preserving all unit distances preserves the distance between x and y to within ε i.e. $||f(x) - f(y)| - |x - y|| \le \varepsilon$.

If $a, b \in C_n$, $a > b > 0$ then $\sqrt{a^2 - b^2} \in C_n$ (see Figure 7).



Figure 7

$|x - y| = \sqrt{a^2 - b^2}$

$S_{xy} = S_{sx} \cup S_{xt} \cup S_{st} \cup S_{sy} \cup S_{ty}$

Hence $\sqrt{3} \cdot a = \sqrt{(2 \cdot a)^2 - a^2} \in C_n$ and $\sqrt{2} \cdot a = \sqrt{(\sqrt{3} \cdot a)^2 - a^2} \in C_n$. Therefore $\sqrt{a^2 + b^2} = \sqrt{(\sqrt{2} \cdot a)^2 - (\sqrt{a^2 - b^2})^2} \in C_n$.

Let us see at Figure 8 below, z denotes the centre of the $(n-1)$-dimensional regular simplex $p_1 p_2 ... p_n$, $n = 2$. This construction shows that if $a, b \in C_n$, $a > b > 0$, $n \geq 2$ (consistently, equations below Figure 8 describe the general case $n \geq 2$) then $a - b \in C_n$, hence $a + b = 2 \cdot a - (a - b) \in C_n$.



Figure 8

$$|x - y| = a - b, \;\; |x - z| = a \in C_n, \;\; |y - z| = b \in C_n$$
$$|p_i - p_j| = \sqrt{2 + 2/n} \in C_n, \;\; |z - p_i| = \sqrt{1^2 - (1/n)^2} \in C_n, \;\; 1 \leq i < j \leq n$$
$$|x - p_1| = \sqrt{|x - z|^2 + |z - p_1|^2} = ... = |x - p_n| = \sqrt{|x - z|^2 + |z - p_n|^2} \in C_n$$
$$|y - p_1| = \sqrt{|y - z|^2 + |z - p_1|^2} = ... = |y - p_n| = \sqrt{|y - z|^2 + |z - p_n|^2} \in C_n$$
$$S_{xy} = \bigcup_{1 \leq i < j \leq n} S_{p_i p_j} \cup \bigcup_{i=1}^{n} S_{x p_i} \cup \bigcup_{i=1}^{n} S_{y p_i} \cup T_{xy}(b)$$

In order to prove that $C_n \backslash \{0\}$ is a multiplicative group it remains to observe that if positive $a, b, c \in C_n$ then $\frac{a \cdot b}{c} \in C_n$ (see Figure 9).



Figure 9

m is positive and integer

$$b < 2 \cdot m \cdot c$$
$$S_{AB} = S_{OA} \cup S_{OB} \cup S_{O\tilde{A}} \cup S_{O\tilde{B}} \cup S_{A\tilde{A}} \cup S_{B\tilde{B}} \cup S_{\tilde{A}\tilde{B}}$$

If $a \in C_n$, $a > 1$, then $\sqrt{a} = \frac{1}{2} \cdot \sqrt{(a+1)^2 - (a-1)^2} \in C_n$; if $a \in C_n$, $0 < a < 1$, then $\sqrt{a} = 1/\sqrt{\frac{1}{a}} \in C_n$. Thus C_n contains all non-negative real numbers contained in the real quadratic closure of Q. This completes the proof.

Directly from the Lemma (see the proof of Theorem 1) we obtain the following two corollaries, these corollaries and the next elementary proposition are useful in our next proofs.

Corollary 1. If $p \neq q \in \mathbf{R}^n$ $(n > 1)$ then any map $f : T_{pq}(|pq|/2) \to \mathbf{R}^n$ that preserves unit distance satisfies $f(p) \neq f(q)$.

Corollary 2. If $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| < |LM|$ then any map $f : T_{JK}(\frac{|LM|-|JK|}{3}) \cup T_{LM}(\frac{|LM|-|JK|}{3}) \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(J)f(K)| < |f(L)f(M)|$.

Proposition. If $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |LM|$ then there exist $A, B \in \mathbf{R}^n$ such that:
(1) segments JK and AB are symmetric with respect to some hyperplane $H(JK, AB)$
and
(2) segments AB and LM are symmetric with respect to some hyperplane $H(AB, LM)$.
Proof. Throughout the proof for any two points $x \in \mathbf{R}^n$ and $y \in \mathbf{R}^n$ H_{xy} stands for the hyperplane which reflects x into y, H_{xy} is unique if $x \neq y$. Let $\widetilde{H}_{xy} : \mathbf{R}^n \to \mathbf{R}^n$ denote the hyperplane symmetry determined by H_{xy}. We put: $A := L = \widetilde{H}_{JL}(J)$, $B := \widetilde{H}_{JL}(K)$, $H(JK, AB) := H_{JL}$. Since $|AB| = |JK| = |LM| = |AM|$ there exists a hyperplane H_{BM} satisfying $\widetilde{H}_{BM}(A) = A = L$. Therefore, we put $H(AB, LM) := H_{BM}$, obviously $\widetilde{H}(AB, LM)(B) = M$. This completes the proof.

Theorems 3 and 4 below yield information about unit-distance preserving mappings on finite subsets of $\mathbf{R}^n$. We present their proofs omitting details.

Theorem 3. If $X_1, X_2, ..., X_m \in \mathbf{R}^n$ $(n > 1)$ lie on some affine hyperplane then there exists a finite set $L(X_1, X_2, ..., X_m) \subseteq \mathbf{R}^n$ containing $X_1, X_2, ..., X_m$ such that each map from $L(X_1, X_2, ..., X_m)$ to $\mathbf{R}^n$ preserving all unit distances preserves the property that $X_1, X_2, ..., X_m$ lie on some affine hyperplane. If $X_1, X_2, ..., X_m \in \mathbf{F}^n$ then we can choose $L(X_1, X_2, ..., X_m) \subseteq \mathbf{F}^n$.

Proof. We choose an affine hyperplane G such that $X_1, X_2, ..., X_m \in G$. We fix $P \in G$ and choose an integer u such that $|PX_i| \leq 2 \cdot u$ for $1 \leq i \leq m$. We fix $O \in \mathbf{R}^n$ such that PO is perpendicular to G and $|PO| = 4 \cdot u$. Let us see at Figure 10 below, this drawing shows the case $n = 2$, but equations below Figure 10 describe the general case $n \geq 2$.



Figure 10

$$\forall i \in \{1, 2, ..., m\} \forall j \in \{1, 2, ..., n\} |X_iB_{ij}| = |A_iB_{ij}| = 2 \cdot u \wedge |OB_{ij}| = 6 \cdot u$$

$$|PO| = 4 \cdot u, \forall i \in \{1, 2, ..., m\} |PA_i| = 4 \cdot u \wedge |PT_i| = |T_iX_i| = u$$

Erasing all dashed lines and segments PT_i, T_iX_i, PA_i, PO from Figure 10 we obtain the Peaucellier inversor: $|OX_i| \cdot |OA_i| = |OC_i| \cdot |OD_i| = (6 \cdot u - 2 \cdot u) \cdot (6 \cdot u + 2 \cdot u) = (6 \cdot u)^2 - (2 \cdot u)^2$ (see [9] in the case $n = 2$). Always: $|PX_i| \leq 2 \cdot u < |PA_i|$. Therefore, if all edge-lengths (continuous segments) are preserved and if all relations $B_{ik} \neq B_{il}$ $(1 \leq k < l \leq n)$ are preserved then PX_i is perpendicular to PO. From this, we conclude that

$$L_i := S_{PO} \cup S_{PT_i} \cup S_{T_iX_i} \cup S_{PA_i} \cup (\bigcup_{j=1}^{n} S_{X_iB_{ij}} \cup S_{A_iB_{ij}} \cup S_{OB_{ij}}) \cup$$
$$\bigcup \{T_{B_{ik}B_{il}}(|B_{ik}B_{il}|/2) : 1 \leq k < l \leq n\}$$

is adequate for preserving the property that PX_i is perpendicular to PO. Hence $L(X_1, X_2, ..., X_m) := \bigcup_{i=1}^{m} L_i$ satisfies the condition of the theorem. This completes the proof for $\mathbf{R}^n$. The proof for $\mathbf{F}^n$ is similar.

Theorem 4 (cf. Corollary 2). If $J, K, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |LM|$ then there exists a finite set $C_{JKLM} \subseteq \mathbf{R}^n$ containing J, K, L, M such that any map $f : C_{JKLM} \to \mathbf{R}^n$ that preserves unit distance satisfies $|f(J)f(K)| = |f(L)f(M)|$. If $J, K, L, M \in \mathbf{F}^n$ then we can choose $C_{JKLM} \subseteq \mathbf{F}^n$.

Proof. Let us observe that if $J, K, A, B, L, M \in \mathbf{R}^n$ $(n > 1)$ and $|JK| = |AB| = |LM|$ then $C_{JKLM} := C_{JKAB} \cup C_{ABLM}$ satisfies the condition of Theorem 4. By Proposition it is sufficient to prove Theorem 4 in the case where two segments of equal lengths are symmetric with respect to some hyperplane H.

Let us see at figures below, drawings show the case $n = 2$, but equations below figures describe the general case $n \geq 2$.



Figure 11a

s, t are positive and integer

$$\forall j \in \{1, 2, ..., n\} |X_jJ| = |X_jL| = s \wedge |Y_jK| = |Y_jM| = t$$
$$|JK| = |LM| < |JM| = |LK| \qquad \delta := \frac{|JM| - |JK|}{3}$$



Figure 11b

s, t are positive and integer

$$\forall j \in \{1, 2, ..., n\} |X_jJ| = |X_jL| = s \wedge |Y_jK| = |Y_jM| = t$$

$$|JK| = |LM| > |JM| = |LK| \qquad \delta := \frac{|JK|-|JM|}{3}$$

If all edge-lengths (continuous segments) are preserved and if all relations: $J \neq L$,$X_k \neq X_l$,$K \neq M$,$Y_k \neq Y_l$ $(1 \leq k < l \leq n)$ are preserved then images of J and L are symmetric and images of K and M are symmetric. Hence: $C_{JKLM} := L(X_1, X_2, ..., X_n, Y_1, Y_2, ..., Y_n) \cup (\bigcup_{j=1}^{n} S_{X_jJ} \cup S_{X_jL}) \cup T_{JL}(|JL|/2) \cup \bigcup\{T_{X_kX_l}(|X_kX_l|/2) : 1 \leq k < l \leq n\} \cup (\bigcup_{j=1}^{n} S_{Y_jK} \cup S_{Y_jM}) \cup T_{KM}(|KM|/2) \cup \bigcup\{T_{Y_kY_l}(|Y_kY_l|/2) : 1 \leq k < l \leq n\} \cup T_{JK}(\delta) \cup T_{LM}(\delta) \cup T_{JM}(\delta) \cup T_{LK}(\delta)$ satisfies the condition of the theorem.

Separate formulas for C_{JKLM} are necessary if $J = L$ or $K = M$. If $J = L$ and $K = M$ then $C_{JKLM} := \{J, K\}$, if $J \neq L$ and $K = M$ then we see at Figure 11c, Figure 11c shows the case $n = 2$, but equations below Figure 11c describe the general case $n \geq 2$.



Figure 11c

s is positive and integer, $\qquad \forall j \in \{1, 2, ..., n\} |X_jJ| = |X_jL| = s$

$$C_{JKLM} = L(X_1, X_2, ..., X_n, K) \cup (\bigcup_{j=1}^{n} S_{X_jJ} \cup S_{X_jL}) \cup T_{JL}(|JL|/2) \cup$$
$$\bigcup\{T_{X_kX_l}(|X_kX_l|/2) : 1 \leq k < l \leq n\}$$

Symmetric construction applies to the case: $J = L$ and $K \neq M$. All cases are explained and the proof for $\mathbf{R}^n$ is complete. The proof for $\mathbf{F}^n$ is similar.

Conjecture. Let us assume that $X \subseteq \underbrace{\mathbf{R}^n \times ... \times \mathbf{R}^n}_{m-times} \cong \mathbf{R}^{mn}$ $(n > 1, m \geq 1)$ is definable using L and the relation X is preserved under isometries of $\mathbf{R}^n$. We suppose that if $(x_1, ..., x_m) \in X$ then there exists a finite set $P(x_1, ..., x_m) \subseteq \mathbf{R}^n$ containing $x_1, ..., x_m$ such that any map $f : P(x_1, ..., x_m) \rightarrow \mathbf{R}^n$ that preserves unit distance satisfies $(f(x_1), ..., f(x_m)) \in X$.

References

1. F. S. Beckman and D. A. Quarles Jr., On isometries of euclidean spaces, *Proc. Amer. Math. Soc.*, 4 (1953), 810-815.
2. W. Benz, *Geometrische Transformationen (unter besonderer Berücksichtigung der Lorentztransformationen).* BI Wissenschaftsverlag, Mannheim, Leipzig, Wien, Zürich, 1992.
3. K. Borsuk, *Multidimensional analytic geometry*, Polish Scientific Publishers, Warsaw, 1969.
4. C. C. Chang and H. J. Keisler, *Model Theory*, 3rd ed., North-Holland, Amsterdam, 1990.
5. U. Everling, Solution of the isometry problem stated by K. Ciesielski, *Math. Intelligencer*, 10 (1988), No.4, p.47.
6. M. Koshelev, We can measure any distance or any amount of time with a most primitive clock and a most primitive ruler: a space-time version of Tyszka's result, *Geombinatorics*, 7 (1998), No.3, 95-100.
7. H. Maehara, Distances in a rigid unit-distance graph in the plane, *Discrete Appl. Math.* 31 (1991), 193-200.
8. H. Maehara, Extending a flexible unit-bar framework to a rigid one, *Discrete Math.* 108 (1992), 167-174.
9. H. Rademacher and O. Toeplitz, *The enjoyment of mathematics*, Princeton University Press, Princeton, 1994.

10. A. Tyszka, A discrete form of the Beckman-Quarles theorem, *Amer. Math. Monthly* 104 (1997), 757-761.

Technical Faculty
Hugo Kołłątaj University
Balicka 104, PL-30-149 Kraków, Poland
rttyszka@cyf-kr.edu.pl
http://www.cyf-kr.edu.pl/~rttyszka